



**2005 TABLES OF
NOTIFIABLE DISEASES**

**Table G. Reported Cases of Selected Notifiable Diseases by Year of Onset
Los Angeles County, 2000-2005**

Disease	Year of Onset						Previous 5-year Average	5-Yr 95% upper Limit ^a
	2000	2001	2002	2003	2004	2005		
Amebiasis	109	139	102	121	114	114	117	142
Botulism ^b	0	2	2	0	3	8	1	4
Brucellosis	4	9	11	7	4	8	7	12
Campylobacteriosis	1273	1141	1067	1100	884	725	1093	1340
Cholera	0	0	0	1	0	0	0	1
Coccidioidomycosis ^b	58	68	76	73	133	214	82	133
Cryptosporidiosis	68	77	62	71	56	45	67	81
Cysticercosis	43	37	18	12	8	15	24	51
Dengue ^b	3	5	7	0	5	10	4	9
<i>E. coli</i> O157:H7	27	31	31	27	18	13	27	36
Encephalitis	49	41	61	38	133	57	64	133
Foodborne outbreaks	40	48	29	25	40	32	36	53
Giardiasis	509	446	441	401	320	313	423	545
<i>Haemophilus influenzae</i> type b	1	5	4	0	2	3	2	6
Hansen's Disease (Leprosy)	9	2	11	9	9	2	8	14
Hepatitis A	839	542	438	374	321	480	503	863
Hepatitis B	65	44	32	73	72	57	57	89
Hepatitis C	10	1	3	0	5	3	4	11
Hepatitis unspecified	11	1	0	1	0	4	3	11
Kawasaki syndrome	35	24	9	14	41	43	25	48
Legionellosis ^b	14	18	25	21	15	31	19	26
Listeriosis, nonperinatal	19	27	14	17	21	25	20	28
Listeriosis, perinatal	8	3	7	3	6	3	5	9
Lyme disease	7	5	8	6	0	7	5	11
Malaria	43	46	38	60	51	45	48	62
Measles	5	8	0	0	1	0	3	9
Meningitis, viral	263	378	466	899	807	515	563	1048
Meningococcal infections	53	58	46	32	28	37	43	66
Mumps	29	17	16	10	5	10	15	31
Pertussis ^b	102	103	172	130	156	438	133	187
Psittacosis	0	1	0	0	0	0	0	1
Q-fever	1	1	4	0	4	0	2	5
Relapsing fever	0	0	1	0	0	0	0	1
Rheumatic fever, acute	1	6	0	0	1	0	2	6
Rubella	3	0	0	0	0	1	1	3
Salmonellosis	990	1006	956	995	1205	1085	1030	1205
Shigellosis	849	684	974	669	625	710	760	1017
Strongyloidiasis	1	0	0	0	0	0	0	1
Tetanus	0	2	2	1	2	0	1	3
Trichinosis	0	0	0	0	0	0	0	0
Tularemia	0	0	0	1	0	0	0	1
Typhoid fever, case	21	17	33	16	13	12	20	34
Typhoid fever, carrier	6	1	6	2	3	4	4	8
Typhus fever	17	8	11	12	8	9	11	18
Vibrio	13	15	14	13	26	14	16	26

^aThe normal distribution assumption may not apply to some rare diseases.

^b2005 data over 95% upper limit.

**Table H. Annual Incidence Rates of Selected Notifiable Diseases by Year of Onset
Los Angeles County, 2000-2005**

Disease	Annual Incidence Rate (Cases per 100,000) ^b					
	2000	2001	2002	2003	2004	2005
Amebiasis	1.22	1.52	1.10	1.29	1.20	1.19
Botulism	-	0.02	0.02	-	0.03	0.08
Brucellosis	0.04	0.10	0.12	0.07	0.04	0.08
Campylobacteriosis	14.19	12.50	11.50	11.70	9.27	7.57
Cholera	-	-	-	0.01	-	-
Coccidioidomycosis	0.65	0.75	0.82	0.78	1.39	2.23
Cryptosporidiosis	0.76	0.84	0.67	0.75	0.59	0.47
Cysticercosis	0.48	0.41	0.19	0.13	0.08	0.16
Dengue	0.03	0.05	0.08	-	0.05	0.10
<i>E. coli</i> O157:H7	0.30	0.34	0.33	0.29	0.19	0.14
Encephalitis	0.55	0.45	0.66	0.40	1.39	0.59
Giardiasis	5.68	4.89	4.75	4.26	3.36	3.27
<i>Haemophilus influenzae</i> type b	0.01	0.05	0.04	-	0.02	0.03
Hansen's Disease (Leprosy)	0.10	0.02	0.12	0.10	0.09	0.02
Hepatitis A	9.36	5.94	4.72	3.98	3.37	5.01
Hepatitis B	0.72	0.48	0.34	0.78	0.76	0.59
Hepatitis C	0.11	0.01	0.03	-	0.05	0.03
Hepatitis unspecified	0.12	0.01	0.00	0.01	-	0.04
Kawasaki syndrome	0.39	0.26	0.10	0.15	0.43	0.45
Legionellosis	0.16	0.20	0.27	0.22	0.16	0.32
Listeriosis, nonperinatal	0.21	0.30	0.15	0.18	0.22	0.26
Listeriosis, perinatal ^a	5.46	2.05	4.96	2.12	4.25	2.14
Lyme disease	0.08	0.05	0.09	0.06	-	0.07
Malaria	0.48	0.50	0.41	0.64	0.53	0.47
Measles	0.06	0.09	-	-	0.01	-
Meningitis, viral	2.93	4.14	5.02	9.56	8.46	5.37
Meningococcal infections	0.59	0.64	0.50	0.34	0.29	0.39
Mumps	0.32	0.19	0.17	0.11	0.05	0.10
Pertussis	1.14	1.13	1.85	1.38	1.64	4.57
Psittacosis	-	0.01	-	-	-	-
Q-fever	0.01	0.01	0.04	-	0.04	-
Relapsing fever	-	-	0.01	-	-	-
Rheumatic fever, acute	0.01	0.07	-	-	0.01	-
Rubella	0.03	-	-	-	-	0.01
Salmonellosis	11.04	11.02	10.30	10.58	12.64	11.33
Shigellosis	9.47	7.50	10.50	7.11	6.55	7.41
Strongyloidiasis	0.01	-	-	-	-	-
Tetanus	-	0.02	0.02	0.01	0.02	-
Trichinosis	-	-	-	-	-	-
Tularemia	-	-	-	0.01	-	-
Typhoid fever, case	0.23	0.19	0.36	0.17	0.14	0.13
Typhoid fever, carrier	0.07	0.01	0.06	0.02	0.03	0.04
Typhus fever	0.19	0.09	0.12	0.13	0.08	0.09
Vibrio	0.14	0.16	0.15	0.14	0.27	0.15

^aRates for perinatal listeriosis were calculated as cases per 100,000 live births.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table I. Five –Year Average
of Notifiable Diseases by Month of Onset
Los Angeles County, 2001-2005**

Disease	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec	Total
Amebiasis	7.2	8.2	9.0	7.6	9.2	8.4	10.0	10.4	9.6	6.8	8.0	9.4	117.8
Botulism	0.2	0.2	0.4	0.2	0.0	0.4	0.2	0.2	0.2	0.0	0.6	0.2	2.8
Brucellosis	0.8	1.0	0.8	0.2	0.2	0.6	1.2	0.8	0.2	1.0	0.6	0.2	7.8
Campylobacteriosis	69.2	53.4	66.2	78.0	94.0	103.6	113.0	102.4	88.6	80.2	76.6	52.2	980.6
Cholera	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.2
Coccidioidomycosis	9.0	5.2	5.6	8.0	8.6	6.6	8.4	11.2	13.0	11.8	10.0	7.6	113.8
Cryptosporidiosis	5.4	3.2	3.4	4.6	4.2	4.0	6.2	9.2	6.4	4.6	4.8	3.8	62.2
Cysticercosis	1.8	1.2	2.6	2.0	3.0	1.0	1.2	1.6	1.0	1.0	0.2	0.4	18.0
Dengue	0.0	0.4	0.0	0.2	0.4	0.4	0.8	1.2	0.8	0.4	0.0	0.0	5.0
E. coli O157:H7	1.4	0.2	1.0	1.0	1.4	2.4	4.8	4.6	3.2	2.2	1.0	0.4	23.8
Encephalitis	3.4	4.0	4.6	3.2	4.4	4.2	7.8	12.4	9.0	4.4	3.2	4.2	68.6
Giardiasis	29.2	21.4	29.0	29.0	29.2	30.2	38.4	38.6	34.0	29.8	26.6	20.0	383.4
<i>Haemophilus influenzae</i> type b	0.6	0.2	0.6	0.0	0.4	0.2	0.0	0.2	0.2	0.2	0.2	0.0	2.8
Hansen's Disease (Leprosy) ^a	-	-	-	-	-	-	-	-	-	-	-	-	-
Hepatitis A	34.8	27.0	28.8	23.8	26.8	23.0	32.0	39.2	42.4	45.8	45.2	37.0	433.2
Hepatitis B	5.8	7.2	5.4	6.0	6.6	6.0	4.2	3.6	2.2	7.0	7.6	7.4	70.0
Hepatitis C	1.0	0.8	0.4	0.8	0.6	1.2	0.4	0.4	0.8	1.0	1.0	2.2	11.4
Hepatitis unspecified	0.0	0.2	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	1.2
Kawasaki syndrome	3.0	2.6	3.2	2.4	2.2	2.8	1.6	2.2	1.4	0.6	1.0	1.8	25.6
Legionellosis	1.2	1.2	1.4	1.4	2.4	2.2	1.6	1.2	0.6	2.2	3.0	1.4	21.8
Listeriosis, nonperinatal	1.0	0.4	1.4	1.4	2.0	3.4	2.2	3.0	1.8	1.4	0.6	1.8	20.2
Listeriosis, perinatal	0.0	0.0	0.4	0.8	0.4	0.4	0.6	0.8	0.6	0.4	0.0	0.0	4.4
Lyme disease	0.2	0.0	0.2	0.0	0.0	0.6	1.0	0.8	0.4	0.4	0.0	0.0	3.6
Malaria ^a	-	-	-	-	-	-	-	-	-	-	-	-	-
Measles	0.2	0.4	0.6	0.2	0.2	0.2	0.0	0.0	0.0	0.0	0.0	0.0	1.8
Meningitis, viral	20.0	16.2	20.0	25.0	29.2	45.6	83.8	109.0	85.4	47.4	37.2	22.6	615.6
Meningococcal infections	6.4	5.4	4.2	5.0	1.8	2.8	2.4	1.6	1.6	2.2	2.2	4.4	40.2
Mumps	0.6	1.2	1.0	0.4	1.2	1.2	1.6	1.2	0.4	1.0	1.0	0.8	11.6
Pertussis	13.6	9.2	11.0	12.6	14.6	16.6	20.4	25.0	23.0	21.0	15.4	17.4	199.8
Psittacosis	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.2
Q-fever	0.4	0.4	0.0	0.2	0.2	0.2	0.0	0.4	0.0	0.0	0.0	0.0	1.8
Relapsing fever	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.2
Rheumatic fever, acute	0.4	0.2	0.4	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.2	0.0	1.4
Rubella	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
Salmonellosis	68.0	50.4	66.4	71.2	96.2	97.8	117.2	117.0	114.0	91.4	69.4	55.0	1044.8
Shigellosis	44.8	29.4	31.8	27.2	33.2	50.8	79.8	114.2	111.0	95.2	56.2	50.2	729.4
Strongyloidiasis	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Tetanus	0.2	0.0	0.0	0.0	0.2	0.2	0.0	0.2	0.0	0.4	0.0	0.2	1.4
Trichinosis	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Tularemia	0.0	0.0	0.0	0.0	0.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.2
Typhoid fever, case	0.8	0.8	1.8	1.6	1.2	2.8	2.0	3.0	2.2	0.8	0.4	0.8	18.2
Typhoid fever, carrier	0.0	0.0	0.8	0.0	0.2	0.6	0.8	0.2	0.2	0.0	0.2	0.2	3.2
Typhus fever	0.4	0.0	0.0	0.2	1.2	1.6	1.0	1.0	1.2	1.0	0.6	0.4	8.6
Vibrio	0.2	0.4	0.6	0.4	0.8	1.4	2.8	2.6	1.8	2.2	1.6	0.4	16.0

^a Not applicable.

**Table J. Number of Cases of Selected Notifiable Diseases by Age Group
Los Angeles County, 2005**

Disease	<1	1-4	5-14	15-34	35-44	45-54	55-64	65+	Total^a
Amebiasis	0	2	14	31	31	26	5	5	114
Botulism	0	0	0	1	4	2	0	1	8
Brucellosis	0	0	2	3	1	0	1	1	8
Campylobacteriosis	31	81	87	203	111	82	56	74	725
Cholera	0	0	0	0	0	0	0	0	0
Coccidioidomycosis	0	0	3	52	50	49	27	33	214
Cryptosporidiosis	0	1	1	10	20	7	4	2	45
Cysticercosis	0	0	0	6	4	1	3	1	15
Dengue	0	0	0	6	2	1	1	0	10
<i>E. coli</i> O157:H7	0	2	4	5	1	1	0	0	13
Encephalitis	3	6	18	11	6	5	0	7	57
Giardiasis	3	37	56	62	58	42	31	23	313
<i>Haemophilus influenzae</i> type b	1	0	1	0	0	0	0	1	3
Hansen's Disease (Leprosy)	0	0	0	1	1	0	0	0	2
Hepatitis A	0	7	24	198	88	88	44	30	480
Hepatitis B	0	0	0	18	21	10	2	6	57
Hepatitis C	0	0	0	1	1	0	1	0	3
Hepatitis unspecified	0	0	0	1	2	0	1	0	4
Kawasaki syndrome	7	29	7	0	0	0	0	0	43
Legionellosis	0	0	0	0	3	5	10	13	31
Listeriosis, nonperinatal	0	2	2	0	0	6	6	9	25
Listeriosis, perinatal ^b	0	0	0	2	1	0	0	0	0
Lyme disease	0	0	1	2	1	1	1	1	7
Malaria	0	0	3	21	8	10	2	1	45
Measles	0	0	0	0	0	0	0	0	0
Meningitis, viral	73	22	91	144	91	46	28	18	515
Meningococcal infections	3	2	6	12	3	3	5	3	37
Mumps	0	1	0	3	0	4	1	1	10
Pertussis	180	27	88	82	32	16	8	5	438
Psittacosis	0	0	0	0	0	0	0	0	0
Q-fever	0	0	0	0	0	0	0	0	0
Relapsing fever	0	0	0	0	0	0	0	0	0
Rheumatic fever, acute	0	0	0	0	0	0	0	0	0
Rubella	1	0	0	0	0	0	0	0	1
Salmonellosis	95	191	189	220	117	88	73	110	1085
Shigellosis	13	170	213	149	70	34	31	28	710
Strongyloidiasis	0	0	0	0	0	0	0	0	0
Tetanus	0	0	0	0	0	0	0	0	0
Trichinosis	0	0	0	0	0	0	0	0	0
Tularemia	0	0	0	0	0	0	0	0	0
Typhoid fever, case	0	1	2	7	0	2	0	0	12
Typhoid fever, carrier	0	0	0	1	0	2	0	1	4
Typhus fever	0	0	1	0	2	3	2	1	9
Vibrio	0	0	1	3	4	3	2	1	14

^aTotals include cases with unknown age.

^bMother's age.

**Table K. Incidence Rates of Selected Notifiable Diseases by Age Group
Los Angeles County, 2005**

Disease	Age-group Rates (Cases per 100,000) ^b							
	<1	1-4	5-14	15-34	35-44	45-54	55-64	65+
Amebiasis	-	0.3	0.9	1.1	2.0	2.0	0.6	0.5
Botulism	-	-	-	-	0.3	0.2	-	0.1
Brucellosis	-	-	0.1	0.1	0.1	-	0.1	0.1
Campylobacteriosis	21.6	14.0	5.9	7.3	7.3	6.4	6.7	7.6
Cholera	-	-	-	-	-	-	-	-
Coccidioidomycosis	-	-	0.2	1.9	3.3	3.8	3.2	3.4
Cryptosporidiosis	-	0.2	0.1	0.4	1.3	0.5	0.5	0.2
Cysticercosis	-	-	-	0.2	0.3	0.1	0.4	0.1
Dengue	-	-	-	0.2	0.1	0.1	0.1	-
<i>E. coli</i> O157:H7	-	0.3	0.3	0.2	0.1	0.1	-	-
Encephalitis	2.1	1.0	1.2	0.4	0.4	0.4	-	0.7
Giardiasis	2.1	6.4	3.8	2.2	3.8	3.3	3.7	2.4
<i>Haemophilus influenzae</i> type b	0.7	-	0.1	-	-	-	-	0.1
Hansen's Disease (Leprosy)	-	-	-	-	0.1	-	-	-
Hepatitis A	-	1.2	1.6	7.1	5.8	6.9	5.2	3.1
Hepatitis B	-	-	-	0.6	1.4	0.8	0.2	0.6
Hepatitis C	-	-	-	-	0.1	-	0.1	-
Hepatitis unspecified	-	-	-	-	0.1	-	0.1	-
Kawasaki syndrome	4.9	5.0	0.5	-	-	-	-	-
Legionellosis	-	-	-	-	0.2	0.4	1.2	1.3
Listeriosis, nonperinatal	-	0.3	0.1	-	-	0.5	0.7	0.9
Listeriosis, perinatal ^a	-	-	1.8	3.9	-	-	-	-
Lyme disease	-	-	0.1	0.1	0.1	0.1	0.1	0.1
Malaria	-	-	0.2	0.8	0.5	0.8	0.2	0.1
Measles	-	-	-	-	-	-	-	-
Meningitis, viral	50.9	3.8	6.2	5.2	6.0	3.6	3.3	1.9
Meningococcal infections	2.1	0.3	0.4	0.4	0.2	0.2	0.6	0.3
Mumps	-	0.2	-	0.1	-	0.3	0.1	0.1
Pertussis	125.6	4.7	6.0	2.9	2.1	1.3	1.0	0.5
Psittacosis	-	-	-	-	-	-	-	-
Q-fever	-	-	-	-	-	-	-	-
Relapsing fever	-	-	-	-	-	-	-	-
Rheumatic fever, acute	-	-	-	-	-	-	-	-
Rubella	0.7	-	-	-	-	-	-	-
Salmonellosis	66.3	33.1	12.8	7.9	7.7	6.9	8.7	11.3
Shigellosis	9.1	29.5	14.4	5.3	4.6	2.7	3.7	2.9
Strongyloidiasis	-	-	-	-	-	-	-	-
Tetanus	-	-	-	-	-	-	-	-
Trichinosis	-	-	-	-	-	-	-	-
Tularemia	-	-	-	-	-	-	-	-
Typhoid fever, case	-	0.2	0.1	0.3	-	0.2	-	-
Typhoid fever, carrier	-	-	-	-	-	0.2	-	0.1
Typhus fever	-	-	0.1	-	0.1	0.2	0.2	0.1
Vibrio	-	-	0.1	0.1	0.3	0.2	0.2	0.1

^aRates for perinatal listeriosis were calculated as cases per 100,000 live births.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table L. Number of Cases of Selected Notifiable Diseases by Race/Ethnicity
Los Angeles County, 2005**

Disease	Asian	Black	Hispanic	White	Other^a	Unknown
Amebiasis	5	7	46	47	2	2
Botulism	2	2	4	0	0	0
Brucellosis	0	0	6	2	0	0
Campylobacteriosis	65	24	318	302	4	11
Cholera	0	0	0	0	0	0
Coccidioidomycosis	15	28	70	96	0	3
Cryptosporidiosis	0	10	16	15	0	4
Cysticercosis	1	1	13	0	0	0
Dengue	4	0	5	1	0	0
<i>E. coli</i> O157:H7	0	0	1	12	0	0
Encephalitis	11	5	27	14	0	0
Giardiasis	20	17	101	149	4	4
<i>Haemophilus influenzae</i> type b	0	0	2	1	0	0
Hansen's Disease (Leprosy)	1	0	1	0	0	0
Hepatitis A	42	49	135	203	13	27
Hepatitis B	8	12	19	16	0	1
Hepatitis C	0	0	0	3	0	0
Hepatitis unspecified	1	0	1	1	0	0
Kawasaki syndrome	14	2	19	6	2	0
Legionellosis	7	2	10	12	0	0
Listeriosis, nonperinatal	5	2	9	9	0	0
Listeriosis, perinatal ^b	0	0	2	1	0	0
Lyme disease	1	0	0	4	0	2
Malaria	7	22	7	6	1	2
Measles	0	0	0	0	0	0
Meningitis, viral	39	55	242	151	3	22
Meningococcal infections	5	2	21	9	0	0
Mumps	4	0	1	4	0	1
Pertussis	14	31	245	147	1	0
Psittacosis	0	0	0	0	0	0
Q-fever	0	0	0	0	0	0
Relapsing fever	0	0	0	0	0	0
Rheumatic fever, acute	0	0	0	0	0	0
Rubella	1	0	0	0	0	0
Salmonellosis	105	74	494	392	7	3
Shigellosis	27	43	500	126	3	9
Strongyloidiasis	0	0	0	0	0	0
Tetanus	0	0	0	0	0	0
Trichinosis	0	0	0	0	0	0
Tularemia	0	0	0	0	0	0
Typhoid fever, case	6	0	6	0	0	0
Typhoid fever, carrier	1	0	3	0	0	0
Typhus fever	0	0	3	6	0	0
Vibrio	1	0	7	4	0	2

^aOther includes Native American and any additional racial group that cannot be categorized as Asian, Black, Hispanic, and White.

^bMother's race.

**Table M. Incidence Rates of Selected Notifiable Diseases by Race/Ethnicity
Los Angeles County, 2005**

Disease	Race/Ethnicity Rates (Cases per 100,000) ^b			
	Asian	Black	Hispanic	White
Amebiasis	0.4	0.8	1.0	1.6
Botulism	0.2	0.2	0.1	-
Brucellosis	-	-	0.1	0.1
Campylobacteriosis	5.2	2.8	7.0	10.5
Cholera	-	-	-	-
Coccidioidomycosis	1.2	3.2	1.5	3.3
Cryptosporidiosis	-	1.2	0.4	0.5
Cysticercosis	0.1	0.1	0.3	-
Dengue	0.3	-	0.1	-
<i>E. coli</i> O157:H7	-	-	-	0.4
Encephalitis	0.9	0.6	0.6	0.5
Giardiasis	1.6	2.0	2.2	5.2
<i>Haemophilus influenzae</i> type b	-	-	-	-
Hansen's Disease (Leprosy)	0.1	-	-	-
Hepatitis A	3.4	5.7	3.0	7.0
Hepatitis B	0.6	1.4	0.4	0.6
Hepatitis C	-	-	-	0.1
Hepatitis unspecified	0.1	-	-	-
Kawasaki syndrome	1.1	0.2	0.4	0.2
Legionellosis	0.6	0.2	0.2	0.4
Listeriosis, nonperinatal	0.4	0.2	0.2	0.3
Listeriosis, perinatal ^a	-	-	2.2	4.1
Lyme disease	0.1	-	-	0.1
Malaria	0.6	2.5	0.2	0.2
Measles	-	-	-	-
Meningitis, viral	3.1	6.4	5.3	5.2
Meningococcal infections	0.4	0.2	0.5	0.3
Mumps	0.3	-	-	0.1
Pertussis	1.1	3.6	5.4	5.1
Psittacosis	-	-	-	-
Q-fever	-	-	-	-
Relapsing fever	-	-	-	-
Rheumatic fever, acute	-	-	-	-
Rubella	0.1	-	-	-
Salmonellosis	8.4	8.6	10.9	13.6
Shigellosis	2.2	5.0	11.0	4.4
Strongyloidiasis	-	-	-	-
Tetanus	-	-	-	-
Trichinosis	-	-	-	-
Tularemia	-	-	-	-
Typhoid fever, case	0.5	-	0.1	-
Typhoid fever, carrier	0.1	-	0.1	-
Typhus fever	-	-	0.1	0.2
Vibrio	0.1	-	0.2	0.1

^aRates for perinatal listeriosis were calculated as cases per 100,000 live births.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table N. Number of Cases and Annual Incidence Rate of Selected Notifiable Diseases by Sex
Los Angeles County, 2005**

Disease	Male		Female	
	Cases	Rate (Cases per 100,000) ^b	Cases	Rate (Cases per 100,000) ^b
Amebiasis	74	1.6	37	0.8
Botulism	6	0.1	2	0.0
Brucellosis	4	0.1	4	0.1
Campylobacteriosis	397	8.4	328	6.8
Cholera	0	-	0	-
Coccidioidomycosis	137	2.9	77	1.6
Cryptosporidiosis	38	0.8	7	0.1
Cysticercosis	9	0.2	6	0.1
Dengue	6	0.1	4	0.1
<i>E. coli</i> O157:H7	9	0.2	4	0.1
Encephalitis	32	0.7	25	0.5
Giardiasis	180	3.8	129	2.7
<i>Haemophilus influenzae</i> type b	2	0.0	1	0.0
Hansen's Disease (Leprosy)	0	-	2	0.0
Hepatitis A	300	6.3	178	3.7
Hepatitis B	45	0.9	12	0.2
Hepatitis C	1	0.0	2	0.0
Hepatitis unspecified	3	0.1	1	0.0
Kawasaki syndrome	28	0.6	15	0.3
Legionellosis	21	0.4	10	0.2
Listeriosis, nonperinatal	13	0.3	12	0.2
Listeriosis, perinatal ^a	1	1.4	-	-
Lyme disease	4	0.1	3	0.1
Malaria	34	0.7	11	0.2
Measles	0	-	0	-
Meningitis, viral	262	5.5	251	5.2
Meningococcal infections	18	0.4	19	0.4
Mumps	3	0.1	7	0.1
Pertussis	201	4.2	237	4.9
Psittacosis	0	-	0	-
Q-fever	0	-	0	-
Relapsing fever	0	-	0	-
Rheumatic fever, acute	0	-	0	-
Rubella	1	0.0	0	-
Salmonellosis	561	11.8	524	10.8
Shigellosis	358	7.5	352	7.3
Strongyloidiasis	0	-	0	-
Tetanus	0	-	0	-
Trichinosis	0	-	0	-
Tularemia	0	-	0	-
Typhoid fever, case	9	0.2	3	0.1
Typhoid fever, carrier	1	0.0	3	0.1
Typhus fever	3	0.1	6	0.1
Vibrio	6	0.1	8	0.2

^aRates for perinatal listeriosis were calculated as cases per 100,000 live births.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table O-1. Selected Notifiable Diseases
SPA 1. Antelope Valley Area
Los Angeles County, 2005**

Disease	Frequency	Rate (Cases per 100,000) ^b
	Antelope	Antelope
Amebiasis	0	-
Botulism	0	-
Brucellosis	0	-
Campylobacteriosis	19	5.5
Cholera	0	-
Coccidioidomycosis	79	23.0
Cryptosporidiosis	0	-
Cysticercosis	0	-
Dengue	1	0.3
<i>E. coli</i> O157:H7	1	0.3
Encephalitis	3	0.9
Giardiasis	9	2.6
<i>Haemophilus influenzae</i> type b	0	-
Hansen's Disease (Leprosy)	1	0.3
Hepatitis A	11	3.2
Hepatitis B	1	0.3
Hepatitis C	0	-
Hepatitis unspecified	0	-
Kawasaki syndrome	2	0.6
Legionellosis	0	-
Listeriosis, nonperinatal	1	0.3
Listeriosis, perinatal ^a	0	-
Lyme disease	0	-
Malaria	2	0.6
Measles	0	-
Meningitis, viral	41	12.0
Meningococcal infections	0	-
Mumps	0	-
Pertussis	46	13.4
Psittacosis	0	-
Q-fever	0	-
Relapsing fever	0	-
Rheumatic fever, acute	0	-
Rubella	0	-
Salmonellosis	28	8.2
Shigellosis	21	6.1
Strongyloidiasis	0	-
Tetanus	0	-
Trichinosis	0	-
Tularemia	0	-
Typhoid fever, case	1	0.3
Typhoid fever, carrier	1	0.3
Typhus fever	0	-
Vibrio	2	0.6

^a Rates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

^b Rates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table O-2. Selected Notifiable Diseases
SPA 2. San Fernando Area
Los Angeles County, 2005**

Disease	Frequency					Rate (Cases per 100,000) ^b				
	EV	GL	SF	WV	TOTAL	EV	GL	SF	WV	TOTAL
Amebiasis	4	10	3	13	30	0.9	2.8	0.6	1.5	1.4
Botulism	1	0	1	0	2	0.2	-	0.2	-	0.1
Brucellosis	0	1	0	2	3	-	0.3	-	0.2	0.1
Campylobacteriosis	30	40	36	95	201	6.7	11.3	7.5	11.1	9.4
Cholera	0	0	0	0	0	-	-	-	-	-
Coccidioidomycosis	6	6	12	52	76	1.3	1.7	2.5	6.1	3.6
Cryptosporidiosis	2	2	0	6	10	0.4	0.6	-	0.7	0.5
Cysticercosis	0	0	0	1	1	-	-	-	0.1	0.0
Dengue	2	1	1	0	4	0.4	0.3	0.2	-	0.2
<i>E. coli</i> O157:H7	0	0	0	1	1	-	-	-	0.1	0.0
Encephalitis	2	3	2	8	15	0.4	0.8	0.4	0.9	0.7
Giardiasis	15	23	14	42	94	3.4	6.5	2.9	4.9	4.4
<i>Haemophilus influenzae</i> type b	0	0	0	1	1	-	-	-	0.1	0.0
Hansen's Disease (Leprosy)	0	0	0	0	0	-	-	-	-	-
Hepatitis A	9	21	11	37	78	2.0	5.9	2.3	4.3	3.7
Hepatitis B	1	2	2	5	10	0.2	0.6	0.4	0.6	0.5
Hepatitis C	0	0	0	1	1	-	-	-	0.1	0.0
Hepatitis unspecified	0	0	0	1	1	-	-	-	0.1	0.0
Kawasaki syndrome	3	1	3	2	9	0.7	0.3	0.6	0.2	0.4
Legionellosis	0	2	1	1	4	-	0.6	0.2	0.1	0.2
Listeriosis, nonperinatal	0	2	0	4	6	-	0.6	-	0.5	0.3
Listeriosis, perinatal ^a	0	0	0	0	0	-	-	-	-	-
Lyme disease	0	1	0	1	2	-	0.3	-	0.1	0.1
Malaria	1	3	4	3	11	0.2	0.8	0.8	0.4	0.5
Measles	0	0	0	0	0	-	-	-	-	-
Meningitis, viral	9	17	31	35	92	2.0	4.8	6.5	4.1	4.3
Meningococcal infections	2	0	3	2	7	0.4	-	0.6	0.2	0.3
Mumps	0	2	0	0	2	-	0.6	-	-	0.1
Pertussis	18	14	28	52	112	4.0	4.0	5.9	6.1	5.3
Psittacosis	0	0	0	0	0	-	-	-	-	-
Q-fever	0	0	0	0	0	-	-	-	-	-
Relapsing fever	0	0	0	0	0	-	-	-	-	-
Rheumatic fever, acute	0	0	0	0	0	-	-	-	-	-
Rubella	0	0	0	1	1	-	-	-	0.1	0.0
Salmonellosis	24	57	47	121	249	5.4	16.1	9.8	14.1	11.7
Shigellosis	19	12	39	63	133	4.3	3.4	8.2	7.4	6.2
Strongyloidiasis	0	0	0	0	0	-	-	-	-	-
Tetanus	0	0	0	0	0	-	-	-	-	-
Trichinosis	0	0	0	0	0	-	-	-	-	-
Tularemia	0	0	0	0	0	-	-	-	-	-
Typhoid fever, case	0	0	1	1	2	-	-	0.2	0.1	0.1
Typhoid fever, carrier	0	0	0	0	0	-	-	-	-	-
Typhus fever	0	1	0	0	1	-	0.3	-	-	0.0
Vibrio	0	0	2	1	3	-	-	0.4	0.1	0.1

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table O-3. Selected Notifiable Diseases
SPA 3. San Gabriel Area
Los Angeles County, 2005**

Disease	Frequency					Rate (Cases per 100,000) ^b				
	AH	EM	FH	PO	TOTAL	AH	EM	FH	PO	TOTAL
Amebiasis	2	2	1	1	6	0.6	0.4	0.3	0.2	0.4
Botulism	1	0	1	0	2	0.3	-	0.3	-	0.1
Brucellosis	0	0	0	1	1	-	-	-	0.2	0.1
Campylobacteriosis	29	12	28	36	105	8.1	2.6	9.0	6.3	6.2
Cholera	0	0	0	0	0	-	-	-	-	-
Coccidioidomycosis	4	1	5	3	13	1.1	0.2	1.6	0.5	0.8
Cryptosporidiosis	1	1	1	1	4	0.3	0.2	0.3	0.2	0.2
Cysticercosis	0	1	0	1	2	-	0.2	-	0.2	0.1
Dengue	0	0	2	0	2	-	-	0.6	-	0.1
<i>E. coli</i> O157:H7	0	0	0	1	1	-	-	-	0.2	0.1
Encephalitis	2	0	2	1	5	0.6	-	0.6	0.2	0.3
Giardiasis	14	1	23	5	43	3.9	0.2	7.4	0.9	2.5
<i>Haemophilus influenzae</i> type b	0	0	0	0	0	-	-	-	-	-
Hansen's Disease (Leprosy)	0	0	0	0	0	-	-	-	-	-
Hepatitis A	18	5	19	14	56	5.1	1.1	6.1	2.5	3.3
Hepatitis B	1	1	0	2	4	0.3	0.2	-	0.4	0.2
Hepatitis C	0	0	0	0	0	-	-	-	-	-
Hepatitis unspecified	1	0	0	0	1	0.3	-	-	-	0.1
Kawasaki syndrome	2	0	1	5	8	0.6	-	0.3	0.9	0.5
Legionellosis	3	0	1	2	6	0.8	-	0.3	0.4	0.4
Listeriosis, nonperinatal	1	0	2	2	5	0.3	-	0.6	0.4	0.3
Listeriosis, perinatal ^a	0	0	0	0	0	-	-	-	-	-
Lyme disease	0	0	0	0	0	-	-	-	-	-
Malaria	2	0	0	3	5	0.6	-	-	0.5	0.3
Measles	0	0	0	0	0	-	-	-	-	-
Meningitis, viral	20	4	35	46	105	5.6	0.9	11.3	8.1	6.2
Meningococcal infections	2	0	2	3	7	0.6	-	0.6	0.5	0.4
Mumps	0	0	0	0	0	-	-	-	-	-
Pertussis	12	2	20	16	50	3.4	0.4	6.4	2.8	2.9
Psittacosis	0	0	0	0	0	-	-	-	-	-
Q-fever	0	0	0	0	0	-	-	-	-	-
Relapsing fever	0	0	0	0	0	-	-	-	-	-
Rheumatic fever, acute	0	0	0	0	0	-	-	-	-	-
Rubella	0	0	0	0	0	-	-	-	-	-
Salmonellosis	49	17	34	61	161	13.7	3.7	10.9	10.7	9.5
Shigellosis	24	10	23	23	80	6.7	2.2	7.4	4.0	4.7
Strongyloidiasis	0	0	0	0	0	-	-	-	-	-
Tetanus	0	0	0	0	0	-	-	-	-	-
Trichinosis	0	0	0	0	0	-	-	-	-	-
Tularemia	0	0	0	0	0	-	-	-	-	-
Typhoid fever, case	0	0	0	0	0	-	-	-	-	-
Typhoid fever, carrier	0	0	0	0	0	-	-	-	-	-
Typhus fever	4	0	1	0	5	1.1	-	0.3	-	0.3
Vibrio	0	0	1	0	1	-	-	0.3	-	0.1

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table O-4. Selected Notifiable Diseases
SPA 4. Metro Area
Los Angeles County, 2005**

Disease	Frequency				Rate (Cases per 100,000) ^b			
	CE	HW	NE	TOTAL	CE	HW	NE	TOTAL
Amebiasis	10	24	3	37	2.7	4.5	0.9	3.0
Botulism	0	0	0	0	-	-	-	-
Brucellosis	0	0	1	1	-	-	0.3	0.1
Campylobacteriosis	22	42	13	77	5.9	7.8	3.9	6.2
Cholera	0	0	0	0	-	-	-	-
Coccidioidomycosis	3	1	6	10	0.8	0.2	1.8	0.8
Cryptosporidiosis	6	11	1	18	1.6	2.0	0.3	1.4
Cysticercosis	0	1	0	1	-	0.2	-	0.1
Dengue	1	0	0	1	0.3	-	-	0.1
<i>E. coli</i> O157:H7	0	1	0	1	-	0.2	-	0.1
Encephalitis	0	4	2	6	-	0.7	0.6	0.5
Giardiasis	11	33	4	48	3.0	6.1	1.2	3.9
<i>Haemophilus influenzae</i> type b	0	0	1	1	-	-	0.3	0.1
Hansen's Disease (Leprosy)	0	0	0	0	-	-	-	-
Hepatitis A	68	42	20	130	18.2	7.8	6.0	10.5
Hepatitis B	5	8	1	14	1.3	1.5	0.3	1.1
Hepatitis C	0	0	0	0	-	-	-	-
Hepatitis unspecified	0	0	0	0	-	-	-	-
Kawasaki syndrome	2	4	3	9	0.5	0.7	0.9	0.7
Legionellosis	0	0	1	1	-	-	0.3	0.1
Listeriosis, nonperinatal	0	0	0	0	-	-	-	-
Listeriosis, perinatal ^a	0	1	1	2	-	1.4	2.0	1.1
Lyme disease	0	1	0	1	-	0.2	-	0.1
Malaria	1	6	1	8	0.3	1.1	0.3	0.6
Measles	0	0	0	0	-	-	-	-
Meningitis, viral	6	13	23	42	1.6	2.4	6.9	3.4
Meningococcal infections	1	6	2	9	0.3	1.1	0.6	0.7
Mumps	0	2	0	2	-	0.4	-	0.2
Pertussis	10	20	7	37	2.7	3.7	2.1	3.0
Psittacosis	0	0	0	0	-	-	-	-
Q-fever	0	0	0	0	-	-	-	-
Relapsing fever	0	0	0	0	-	-	-	-
Rheumatic fever, acute	0	0	0	0	-	-	-	-
Rubella	0	0	0	0	-	-	-	-
Salmonellosis	32	83	33	148	8.6	15.5	9.9	11.9
Shigellosis	52	62	32	146	14.0	11.5	9.6	11.7
Strongyloidiasis	0	0	0	0	-	-	-	-
Tetanus	0	0	0	0	-	-	-	-
Trichinosis	0	0	0	0	-	-	-	-
Tularemia	0	0	0	0	-	-	-	-
Typhoid fever, case	0	0	0	0	-	-	-	-
Typhoid fever, carrier	0	0	0	0	-	-	-	-
Typhus fever	1	2	0	3	0.3	0.4	-	0.2
Vibrio	0	1	0	1	-	0.2	-	0.1

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table O-5. Selected Notifiable Diseases
SPA 5. West Area
Los Angeles County, 2005**

Disease	Frequency	Rate (Cases per 100,000) ^b
	West	West
Amebiasis	17	2.6
Botulism	0	-
Brucellosis	0	-
Campylobacteriosis	107	16.5
Cholera	0	-
Coccidioidomycosis	4	0.6
Cryptosporidiosis	3	0.5
Cysticercosis	0	-
Dengue	0	-
<i>E. coli</i> O157:H7	2	0.3
Encephalitis	1	0.2
Giardiasis	34	5.2
<i>Haemophilus influenzae</i> type b	1	0.2
Hansen's Disease (Leprosy)	0	-
Hepatitis A	45	6.9
Hepatitis B	5	0.8
Hepatitis C	2	0.3
Hepatitis unspecified	0	-
Kawasaki syndrome	1	0.2
Legionellosis	1	0.2
Listeriosis, nonperinatal	4	0.6
Listeriosis, perinatal ^a	0	-
Lyme disease	2	0.3
Malaria	3	0.5
Measles	0	-
Meningitis, viral	11	1.7
Meningococcal infections	0	-
Mumps	5	0.8
Pertussis	31	4.8
Psittacosis	0	-
Q-fever	0	-
Relapsing fever	0	-
Rheumatic fever, acute	0	-
Rubella	0	-
Salmonellosis	87	13.4
Shigellosis	43	6.6
Strongyloidiasis	0	-
Tetanus	0	-
Trichinosis	0	-
Tularemia	0	-
Typhoid fever, case	1	0.2
Typhoid fever, carrier	0	-
Typhus fever	0	-
Vibrio	3	0.5

^a Rates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

^b Rates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table O-6. Selected Notifiable Diseases
SPA 6. South Area
Los Angeles County, 2005**

Disease	Frequency					Rate (Cases per 100,000) ^b				
	CN	SO	SE	SW	TOTAL	CN	SO	SE	SW	TOTAL
Amebiasis	0	4	2	3	9	-	2.2	1.2	0.8	0.9
Botulism	0	1	0	0	1	-	0.6	-	-	0.1
Brucellosis	0	1	1	0	2	-	0.6	0.6	-	0.2
Campylobacteriosis	20	5	15	14	54	6.9	2.8	8.9	3.5	5.2
Cholera	0	0	0	0	0	-	-	-	-	-
Coccidioidomycosis	2	2	2	4	10	0.7	1.1	1.2	1.0	1.0
Cryptosporidiosis	0	2	1	1	4	-	1.1	0.6	0.3	0.4
Cysticercosis	0	0	0	2	2	-	-	-	0.5	0.2
Dengue	0	0	0	1	1	-	-	-	0.3	0.1
<i>E. coli</i> O157:H7	0	0	0	0	0	-	-	-	-	-
Encephalitis	0	0	3	0	3	-	-	1.8	-	0.3
Giardiasis	2	5	3	13	23	0.7	2.8	1.8	3.3	2.2
<i>Haemophilus influenzae</i> type b	0	0	0	0	0	-	-	-	-	-
Hansen's Disease (Leprosy)	0	0	0	0	0	-	-	-	-	-
Hepatitis A	7	4	11	8	30	2.4	2.2	6.5	2.0	2.9
Hepatitis B	5	0	0	2	7	1.7	-	-	0.5	0.7
Hepatitis C	0	0	0	0	0	-	-	-	-	-
Hepatitis unspecified	0	0	0	0	0	-	-	-	-	-
Kawasaki syndrome	0	1	0	2	3	-	0.6	-	0.5	0.3
Legionellosis	1	0	0	1	2	0.3	-	-	0.3	0.2
Listeriosis, nonperinatal	0	1	1	1	3	-	0.6	0.6	0.3	0.3
Listeriosis, perinatal ^a	0	0	1	0	1	-	-	3.6	-	0.6
Lyme disease	0	0	0	0	0	-	-	-	-	-
Malaria	1	2	1	3	7	0.3	1.1	0.6	0.8	0.7
Measles	0	0	0	0	0	-	-	-	-	-
Meningitis, viral	24	5	2	7	38	8.3	2.8	1.2	1.8	3.7
Meningococcal infections	1	1	2	1	5	0.3	0.6	1.2	0.3	0.5
Mumps	0	0	0	0	0	-	-	-	-	-
Pertussis	19	11	13	18	61	6.6	6.1	7.7	4.5	5.9
Psittacosis	0	0	0	0	0	-	-	-	-	-
Q-fever	0	0	0	0	0	-	-	-	-	-
Relapsing fever	0	0	0	0	0	-	-	-	-	-
Rheumatic fever, acute	0	0	0	0	0	-	-	-	-	-
Rubella	0	0	0	0	0	-	-	-	-	-
Salmonellosis	23	26	16	44	109	8.0	14.4	9.5	11.1	10.5
Shigellosis	24	24	46	26	120	8.3	13.3	27.2	6.5	11.6
Strongyloidiasis	0	0	0	0	0	-	-	-	-	-
Tetanus	0	0	0	0	0	-	-	-	-	-
Trichinosis	0	0	0	0	0	-	-	-	-	-
Tularemia	0	0	0	0	0	-	-	-	-	-
Typhoid fever, case	0	0	1	2	3	-	-	0.6	0.5	0.3
Typhoid fever, carrier	0	0	0	1	1	-	-	-	0.3	0.1
Typhus fever	0	0	0	0	0	-	-	-	-	-
Vibrio	1	0	0	1	2	0.3	-	-	0.3	0.2

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table O-7. Selected Notifiable Diseases
SPA 7. East Area
Los Angeles County, 2005**

Disease	Frequency					Rate (Cases per 100,000) ^b				
	BF	EL	SA	WH	TOTAL	BF	EL	SA	WH	TOTAL
Amebiasis	1	2	4	2	9	0.3	0.9	0.9	0.6	0.7
Botulism	0	0	0	0	0	-	-	-	-	-
Brucellosis	0	0	0	0	0	-	-	-	-	-
Campylobacteriosis	18	14	25	24	81	4.9	6.2	5.6	7.4	5.9
Cholera	0	0	0	0	0	-	-	-	-	-
Coccidioidomycosis	4	2	7	3	16	1.1	0.9	1.6	0.9	1.2
Cryptosporidiosis	0	2	0	2	4	-	0.9	-	0.6	0.3
Cysticercosis	2	0	4	0	6	0.5	-	0.9	-	0.4
Dengue	0	0	0	0	0	-	-	-	-	-
<i>E. coli</i> O157:H7	0	0	0	2	2	-	-	-	0.6	0.1
Encephalitis	5	2	2	1	10	1.4	0.9	0.4	0.3	0.7
Giardiasis	9	3	11	7	30	2.4	1.3	2.5	2.1	2.2
<i>Haemophilus influenzae</i> type b	0	0	0	0	0	-	-	-	-	-
Hansen's Disease (Leprosy)	1	0	0	0	1	0.3	-	-	-	0.1
Hepatitis A	12	12	14	12	50	3.2	5.3	3.1	3.7	3.6
Hepatitis B	4	2	0	2	8	1.1	0.9	-	0.6	0.6
Hepatitis C	0	0	0	0	0	-	-	-	-	-
Hepatitis unspecified	0	0	0	0	0	-	-	-	-	-
Kawasaki syndrome	1	0	3	0	4	0.3	-	0.7	-	0.3
Legionellosis	0	2	0	4	6	-	0.9	-	1.2	0.4
Listeriosis, nonperinatal	1	1	0	1	3	0.3	0.4	-	0.3	0.2
Listeriosis, perinatal ^a	0	0	0	0	0	-	-	-	-	-
Lyme disease	0	0	0	0	0	-	-	-	-	-
Malaria	1	0	2	0	3	0.3	-	0.4	-	0.2
Measles	0	0	0	0	0	-	-	-	-	-
Meningitis, viral	38	11	26	39	114	10.3	4.8	5.8	12.0	8.3
Meningococcal infections	1	0	3	2	6	0.3	-	0.7	0.6	0.4
Mumps	0	0	0	0	0	-	-	-	-	-
Pertussis	6	8	18	7	39	1.6	3.5	4.0	2.1	2.8
Psittacosis	0	0	0	0	0	-	-	-	-	-
Q-fever	0	0	0	0	0	-	-	-	-	-
Relapsing fever	0	0	0	0	0	-	-	-	-	-
Rheumatic fever, acute	0	0	0	0	0	-	-	-	-	-
Rubella	0	0	0	0	0	-	-	-	-	-
Salmonellosis	49	31	56	21	157	13.2	13.6	12.5	6.4	11.4
Shigellosis	17	21	49	20	107	4.6	9.2	10.9	6.1	7.8
Strongyloidiasis	0	0	0	0	0	-	-	-	-	-
Tetanus	0	0	0	0	0	-	-	-	-	-
Trichinosis	0	0	0	0	0	-	-	-	-	-
Tularemia	0	0	0	0	0	-	-	-	-	-
Typhoid fever, case	0	2	0	0	2	-	0.9	-	-	0.1
Typhoid fever, carrier	0	2	0	0	2	-	0.9	-	-	0.1
Typhus fever	0	0	0	0	0	-	-	-	-	-
Vibrio	0	0	0	1	1	-	-	-	0.3	0.1

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.

**Table O-8. Selected Notifiable Diseases
SPA 8. South Bay Area
Los Angeles County, 2005**

Disease	Frequency				Rate (Cases per 100,000) ^b			
	HB	IW	TO	TOTAL	HB	IW	TO	TOTAL
Amebiasis	1	3	2	6	0.5	0.7	0.4	0.5
Botulism	1	2	0	3	0.5	0.5	-	0.3
Brucellosis	0	1	0	1	-	0.2	-	0.1
Campylobacteriosis	19	34	28	81	9.1	7.9	6.0	7.3
Cholera	0	0	0	0	-	-	-	-
Coccidioidomycosis	1	0	4	5	0.5	-	0.9	0.5
Cryptosporidiosis	0	1	1	2	-	0.2	0.2	0.2
Cysticercosis	1	1	1	3	0.5	0.2	0.2	0.3
Dengue	1	0	0	1	0.5	-	-	0.1
<i>E. coli</i> O157:H7	3	0	2	5	1.4	-	0.4	0.5
Encephalitis	8	1	1	10	3.8	0.2	0.2	0.9
Giardiasis	8	8	16	32	3.8	1.9	3.4	2.9
<i>Haemophilus influenzae</i> type b	0	0	0	0	-	-	-	-
Hansen's Disease (Leprosy)	0	0	0	0	-	-	-	-
Hepatitis A	11	24	23	58	5.3	5.6	4.9	5.2
Hepatitis B	2	2	4	8	1.0	0.5	0.9	0.7
Hepatitis C	0	0	0	0	-	-	-	-
Hepatitis unspecified	0	0	0	0	-	-	-	-
Kawasaki syndrome	2	3	2	7	1.0	0.7	0.4	0.6
Legionellosis	0	0	1	1	-	-	0.2	0.1
Listeriosis, nonperinatal	0	1	1	2	-	0.2	0.2	0.2
Listeriosis, perinatal ^a	0	0	0	0	-	-	-	-
Lyme disease	1	0	1	2	0.5	-	0.2	0.2
Malaria	1	3	2	6	0.5	0.7	0.4	0.5
Measles	0	0	0	0	-	-	-	-
Meningitis, viral	13	31	26	70	6.2	7.2	5.6	6.3
Meningococcal infections	0	1	2	3	-	0.2	0.4	0.3
Mumps	0	0	1	1	-	-	0.2	0.1
Pertussis	20	21	21	62	9.6	4.9	4.5	5.6
Psittacosis	0	0	0	0	-	-	-	-
Q-fever	0	0	0	0	-	-	-	-
Relapsing fever	0	0	0	0	-	-	-	-
Rheumatic fever, acute	0	0	0	0	-	-	-	-
Rubella	0	0	0	0	-	-	-	-
Salmonellosis	55	40	46	141	26.4	9.3	9.8	12.8
Shigellosis	14	21	25	60	6.7	4.9	5.3	5.4
Strongyloidiasis	0	0	0	0	-	-	-	-
Tetanus	0	0	0	0	-	-	-	-
Trichinosis	0	0	0	0	-	-	-	-
Tularemia	0	0	0	0	-	-	-	-
Typhoid fever, case	1	2	0	3	0.5	0.5	-	0.3
Typhoid fever, carrier	0	0	0	0	-	-	-	-
Typhus fever	0	0	0	0	-	-	-	-
Vibrio	0	0	1	1	-	-	0.2	0.1

^aRates for perinatal listeriosis were calculated as cases per 100,000 women aged 15 to 44 years.

^bRates of disease based on less than 20 cases or events are considered "unreliable." A zero rate made from no events is especially hazardous and are not reported here, except with a dash ("-"). Conclusions drawn from unreliable rates should be made with caution, if they are to be made at all.