

How to take DoxyPEP

(*Doxycycline Post-Exposure Prophylaxis*)

DoxyPEP: Protects against STIs such as Chlamydia, Syphilis and Gonorrhea

Two 100 mg pills of doxycycline ideally **within 24 hours** but **no later than 72 hours** after the last time you had condomless oral, anal or vaginal sex.


Example: Sex on Sunday; take dose by Wednesday.



Reminder:
No more than
200mg per 24
hours.



Example: Sex Sunday through Tuesday; take initial dose within 24 hours (by Monday), and daily doses through Tuesday. Last dose should be no later than 72 hours after last sexual encounter (by Friday).



How to take HIV PrEP

(HIV Pre-Exposure Prophylaxis)

HIV PrEP: Protects against HIV

	Oral	2-1-1 (On Demand)	Injectable
Formulation	Tenofovir disoproxil fumarate/emtricitabine (TDF/FTC) [Truvada®] OR Tenofovir alafenamide/emtricitabine (TAF/FTC) [Descovy®]	Tenofovir disoproxil fumarate/emtricitabine (TDF/FTC) [Truvada®]	Cabotegravir (CAB-LA) [Apretude®]
Dosing	One pill daily	Take 2 tablets 2-24 hours before sex, 1 tablet 24 hours after first dose, and another tablet 24 hours later	One injection every 2 months after two initiation injections administered 1 month apart

DoxyPEP is safe to be taken with HIV PrEP!



Ending
the
HIV
Epidemic





DoxyPEP

(Doxycycline Post-Exposure Prophylaxis)

DoxyPEP is **delivered in a pill form**, and can be taken when possible exposure to an STD after sex without barriers.

DoxyPEP must be taken ideally **24 hours after possible sexual exposure**; **no later than 72 hours after possible sexual exposure**.



HIV PrEP

(HIV Pre-Exposure Prophylaxis)



HIV PrEP is **delivered in a pill form OR as an injectable**.

HIV PrEP **must be orally taken daily OR as an injection every 2 months** to prevent HIV acquisition.

