




Selected Maternal Health Indicators for Asian/Pacific Islanders 2005 Los Angeles Mommy and Baby (LAMB) Project

Self-Reported Maternal Medical Conditions During Pregnancy



Preconception Health Indicators		
	Asian/Pacific Islanders	LA County
Uninsured	16.8%	35.6%
Unintended pregnancy	32.4%	40.0%
Did not Meet MediCal Prenatal Care Guidelines	%	17.0%
Late onset of prenatal care	%	7.2%
Did not take multi-vitamin	%	57.0%
Smoking six months prior to becoming pregnant	8.4%	9.2%
Prior Obstetrical history		
Prior miscarriage	19.4%	16.6%
Preterm or low birth weight	5.8%	9.4%
Fetal Death	1.6%	1.8%
Maternal medical condition prior to pregnancy:		
Anemia	7.8%	9.8%
Periodontal disease	4.6%	5.4%
High blood pressure	1.6%	2.6%
Diabetes	2.6%	3.0%

Psychosocial and Risk Behavior Indicators

	Asian/Pacific Islanders	LA County
Depressed during pregnancy	21.0%	34.0%
Lost interest in hobbies or things that they normally enjoyed	20.8%	28.8%
Smoking during pregnancy	2.2%	2.8%
Exposed to second hand smoke	8.4%	10.8%
Drinking during pregnancy	11.0%	12.6%
Using drugs during pregnancy	2.6%	5.4%

What is LAMB Project?

- A population-based survey of newly delivered moms stratified by SPA and race/ethnicity from birth records
- Close to 5,500 women responded to the survey in 2005 LAMB, of which about 1075 were Asian/Pacific Islander women
- Mail survey with telephone followup administered in three languages
- Adjusted response rate was close to 50%


GLOSSARY OF MATERNAL/INFANT HEALTH INDICATOR TERMS

- 1) Depressed during pregnancy: defined as for two weeks or longer during pregnancy, feeling depressed for most of the day or losing interest in work, hobbies, or things usually enjoyed.
- 2) Using drugs during pregnancy: defined as using any drugs that are not prescribed by a physician
- 3) Unintended pregnancy: defined as, just before becoming pregnant, wanting to be pregnant later (i.e. mistimed) or not wanting to be pregnant then or at any time in the future (i.e. unwanted).
- 4) Meeting the medical prenatal care recommended guidelines: defined as meeting all the recommendations of the Public Health Service Expert Panel on the Content of Prenatal Care (1989), including having blood pressure measured, urine and blood samples taken, height and weight measured, a pelvic exam, and a health history taken.
- 5) Late onset of prenatal care: defined as having started prenatal care after the first trimester.



Selected Maternal Health Indicators for African American 2005 Los Angeles Mommy and Baby (LAMB) Project

Self-Reported Maternal Medical Conditions During Pregnancy



Preconception Health Indicators

	African American	LA County
Uninsured	16.4%	35.6%
Unintended pregnancy	56.2%	40.0%
Did not Meet MediCal Prenatal Care Guidelines	14.2%	17.0%
Late onset of prenatal care	12.6%	7.2%
Did not take multi-vitamin	57.4%	57.0%
Smoking six months prior to becoming pregnant	19.4%	9.2%
Prior Obstetrical history		
Prior miscarriage	28.2%	16.6%
Preterm or low birth weight	11.8%	9.4%
Fetal Death	4.4%	1.8%
Maternal medical condition prior to pregnancy:		
Anemia	22.4%	9.8%
Periodontal disease	2.8%	5.4%
High blood pressure	5.6%	2.6%
Diabetes	1.8%	3.0%

Psychosocial and Risk Behavior Indicators

	African American	LA County
Depressed during pregnancy	38.2%	34.0%
Lost interest in hobbies or things that they normally enjoyed	37.6%	28.8%
Smoking during pregnancy	10.8%	2.8%
Exposed to second hand smoke	19.8%	10.8%
Drinking during pregnancy	8.8%	12.6%
Using drugs during pregnancy	7.8%	5.4%

What is LAMB Project?

- A population-based survey of newly delivered moms stratified by SPA and race/ethnicity from birth records
- Close to 5,500 women responded to the survey in 2005 LAMB, of which about 1100 were African American women
- Mail survey with telephone followup administered in three languages
- Adjusted response rate was close to 50%


GLOSSARY OF MATERNAL/INFANT HEALTH INDICATOR TERMS

- 1) Depressed during pregnancy: defined as for two weeks or longer during pregnancy, feeling depressed for most of the day or losing interest in work, hobbies, or things usually enjoyed.
- 2) Using drugs during pregnancy: defined as using any drugs that are not prescribed by a physician
- 3) Unintended pregnancy: defined as, just before becoming pregnant, wanting to be pregnant later (i.e. mistimed) or not wanting to be pregnant then or at any time in the future (i.e. unwanted).
- 4) Meeting the medical prenatal care recommended guidelines: defined as meeting all the recommendations of the Public Health Service Expert Panel on the Content of Prenatal Care (1989), including having blood pressure measured, urine and blood samples taken, height and weight measured, a pelvic exam, and a health history taken.
- 5) Late onset of prenatal care: defined as having started prenatal care after the first trimester.



Selected Maternal Health Indicators for Hispanic 2005 Los Angeles Mommy and Baby (LAMB) Project

Self-Reported Maternal Medical Conditions During Pregnancy



Psychosocial and Risk Behavior Indicators

	Hispanic	LA County
Depressed during pregnancy	39.6%	34.0%
Lost interest in hobbies or things that they normally enjoyed	32.6%	28.8%
Smoking during pregnancy	1.8%	2.8%
Exposed to second hand smoke	11.0%	10.8%
Drinking during pregnancy	8.6%	12.6%
Using drugs during pregnancy	5.6%	5.4%

What is LAMB Project?

- A population-based survey of newly delivered moms stratified by SPA and race/ethnicity from birth records
- Close to 5,500 women responded to the survey in 2005 LAMB, of which about 1630 were Hispanic women
- Mail survey with telephone followup administered in three languages
- Adjusted response rate was close to 50%

Preconception Health Indicators

	Hispanic	LA County
Uninsured	47.2%	35.6%
Unintended pregnancy	44.4%	40.0%
Did not Meet MediCal Prenatal Care Guidelines	16.8%	17.0%
Late onset of prenatal care	7.4%	7.2%
Did not take multi-vitamin	65.6%	57.0%
Smoking six months prior to becoming pregnant	6.8%	9.2%
Prior Obstetrical history		
Prior miscarriage	12.0%	16.6%
Preterm or low birth weight	10.4%	9.4%
Fetal Death	1.4%	1.8%
Maternal medical condition prior to pregnancy:		
Anemia	9.8%	9.8%
Periodontal disease	6.2%	5.4%
High blood pressure	2.8%	2.6%
Diabetes	4.0%	3.0%


GLOSSARY OF MATERNAL/INFANT HEALTH INDICATOR TERMS

- 1) Depressed during pregnancy: defined as for two weeks or longer during pregnancy, feeling depressed for most of the day or losing interest in work, hobbies, or things usually enjoyed.
- 2) Using drugs during pregnancy: defined as using any drugs that are not prescribed by a physician
- 3) Unintended pregnancy: defined as, just before becoming pregnant, wanting to be pregnant later (i.e. mistimed) or not wanting to be pregnant then or at any time in the future (i.e. unwanted).
- 4) Meeting the medical prenatal care recommended guidelines: defined as meeting all the recommendations of the Public Health Service Expert Panel on the Content of Prenatal Care (1989), including having blood pressure measured, urine and blood samples taken, height and weight measured, a pelvic exam, and a health history taken.
- 5) Late onset of prenatal care: defined as having started prenatal care after the first trimester.



Selected Maternal Health Indicators for White 2005 Los Angeles Mommy and Baby (LAMB) Project

Self-Reported Maternal Medical Conditions During Pregnancy



Psychosocial and Risk Behavior Indicators

	White	LA County
Depressed during pregnancy	18.4%	34.0%
Lost interest in hobbies or things that they normally enjoyed	16.6%	28.8%
Smoking during pregnancy	4.2%	2.8%
Exposed to second hand smoke	8.6%	10.8%
Drinking during pregnancy	29.2%	12.6%
Using drugs during pregnancy	5.2%	5.4%

What is LAMB Project?

- A population-based survey of newly delivered moms stratified by SPA and race/ethnicity from birth records
- Close to 5,500 women responded to the survey in 2005 LAMB, of which about 1330 were White women
- Mail survey with telephone followup administered in three languages
- Adjusted response rate was close to 50%

Preconception Health Indicators

	White	LA County
Uninsured	9.6%	35.6%
Unintended pregnancy	22.4%	40.0%
Did not Meet MediCal Prenatal Care Guidelines	19.4%	17.0%
Late onset of prenatal care	5.4%	7.2%
Did not take multi-vitamin	33.2%	57.0%
Smoking six months prior to becoming pregnant	15.2%	9.2%
Prior Obstetrical history		
Prior miscarriage	26.6%	16.6%
Preterm or low birth weight	7.4%	9.4%
Fetal Death	1.8%	1.8%
Maternal medical condition prior to pregnancy:		
Anemia	6.8%	9.8%
Periodontal disease	3.4%	5.4%
High blood pressure	1.8%	2.6%
Diabetes	0.4%	3.0%

GLOSSARY OF MATERNAL/INFANT HEALTH INDICATOR TERMS

- 1) Depressed during pregnancy: defined as for two weeks or longer during pregnancy, feeling depressed for most of the day or losing interest in work, hobbies, or things usually enjoyed.
- 2) Using drugs during pregnancy: defined as using any drugs that are not prescribed by a physician
- 3) Unintended pregnancy: defined as, just before becoming pregnant, wanting to be pregnant later (i.e. mistimed) or not wanting to be pregnant then or at any time in the future (i.e. unwanted).
- 4) Meeting the medical prenatal care recommended guidelines: defined as meeting all the recommendations of the Public Health Service Expert Panel on the Content of Prenatal Care (1989), including having blood pressure measured, urine and blood samples taken, height and weight measured, a pelvic exam, and a health history taken.
- 5) Late onset of prenatal care: defined as having started prenatal care after the first trimester.