

Epidemiology of Tuberculosis in Los Angeles County

Alicia H. Chang, MD MS
Acting Director
Los Angeles County TB Control Program

April 1, 2016




Learning Objectives


- Describe the basic trends and patterns of tuberculosis in Los Angeles County in 2015
- Describe the epidemiology of TB among specific population subsets
- Describe the demographics and clinical characteristics of our TB cases
- Describe two current tuberculosis outbreaks in LA County



TB Cases in LA County, 1992-2015



TB patients by Service Planning Area 2015




DEMOGRAPHICS of TUBERCULOSIS PATIENTS




TB Case Rates by Sex: LA County 2001-2015

Female Male



Age-Gender Distributions: LA County 2014-2015




LA County TB patients, 2015

LA County population (2014)




LA County TB Patients. Data exclude Pasadena and Long Beach TB patients. Based on TRIMS data, updated 2/8/16. Data are provisional and subject to change.


TB Case Rates by Age: LA County 2011-2015




TB Case Rates by Sex and Age: LA County 2015



Tuberculosis in children under 5 years: LA County 2001-2015




TB Rates by Race/Ethnicity: LA County 2001-2015




**Asian includes NHPI (Native Hawaiian / Pacific Islander). LA County TB Patients. Data exclude Pasadena and Long Beach TB patients. Based on TRIMS data, updated 2/8/16. Data are provisional and subject to change. Population estimates source: Internal Services Department Social Services Systems Division, 2014 Population Estimates, prepared for County of Los Angeles, 2014.. 2015 population estimates based on latest available data (2014).


U.S. and Non-U.S. Born TB Patients: LA County 1992-2015




U.S. and Non-U.S. Born TB Patients by Race/Ethnicity: LA County 2015



TB Patients by Nativity and Race/Ethnicity: LA County 2015




TB Patients by Nativity and Country of Birth: LA County 2015




TB Patients by Country of Birth (8 highest): LA County 2011-2015

2011 2012 2013 2014 2015



*Korea includes Democratic People's Republic of Korea & Republic of Korea
LA County TB Patients. Data exclude Pasadena and Long Beach TB patients.
Based on TRIMS data, updated 2/8/16. Data are provisional and subject to change.


Non-U.S. Born Patients by Years in the U.S. at TB diagnosis: LA County 2011-2015




LA County TB Patients. Data exclude Pasadena and Long Beach TB patients. Based on TRIMS data, updated 2/8/16. Data are provisional and subject to change.




Tuberculosis Patients with known HIV Co-Infection: LA County 2001-2015



TB Patients missing HIV Status: LA County 2001-2015




Key Comorbidities among TB Patients: LA County 2011-2015



NOTE: Patients may have more than one comorbidity




Number of Comorbidities* among TB Patients: LA County 2015 (n=606)



*Includes HIV, DM, Immunosuppressed, ESRD, Organ transplant, TNF-alpha therapy




TB Patients by Behavioral Risk Factors: LA County 2010-2015



NOTE: Patients may have more than one risk factor



TB Patients by Social Risk Factors: LA County 2010-2015




NOTE: Patients may have more than one risk factor




CLINICAL CHARACTERISTICS of TUBERCULOSIS PATIENTS



Primary Reason for TB Evaluation: LA County 2015



TB Patients by Site of Disease: LA County 2015 (N=606)



TB Patients by Smear Positivity and Type of Confirmation: LA County 2015

Site of TB Disease	Sputum Smear Positive		Sputum Culture Positive**	Other Culture Positive**	NAAT Positive	Clinical Confirmation
	No	Yes				
	%	%	%	%	%	%
Pulmonary	37.5	62.5	88.3	70.2	72.7	7.6
Extra-pulmonary	98.4	1.6	0	75	16.7	17.4
Both Pulmonary and Extra-pulmonary	54.8	45.2	78.7	79.3	63	4.5

*Smear and culture positivity defined as within 14 days of treatment start date. **Sputum culture includes 'sputum' and 'direct sputum' specimens only. Other culture excludes 'sputum' and 'direct sputum' specimens. Data exclude Long Beach and Pasadena TB cases. Data are provisional and subject to change.




INH mono-resistance

Country	Total TB Cases	Culture positive cases with DST	INH resistance		MDR	
			Yes	%INH resistant	Yes	%MDR
	Number of Cases	Number of Cases	Number of Cases	%INH resistant	Number of Cases	%MDR
Mexico	2096	1624	126	7.8%	15	0.9%
United States	1747	1221	56	4.6%	7	0.6%
Philippines	1191	1015	161	15.9%	18	1.8%
China	447	371	32	8.6%	9	2.4%
Vietnam	429	364	60	16.5%	4	1.1%
Korea*	402	334	51	15.3%	11	3.3%
Guatemala	346	283	27	9.5%	5	1.8%
El Salvador	293	223	14	6.3%	1	0.4%
India	153	122	11	9.0%	2	1.6%
Taiwan	87	76	6	7.9%	0	0.0%
Honduras	78	67	3	4.5%	1	1.5%
Peru	71	61	7	11.5%	3	4.9%
Cambodia	72	58	9	15.5%	3	5.2%
Iran	68	57	4	7.0%	0	0.0%
Indonesia	61	54	6	11.1%	1	1.9%
Armenia	55	46	3	6.5%	2	4.3%
Ethiopia	50	41	3	7.3%	1	2.4%
Burma	43	34	2	5.9%	1	2.9%
Thailand	45	34	4	11.8%	2	5.9%
Japan	35	32	0	0.0%	0	0.0%
Hong Kong	30	26	2	7.7%	0	0.0%




Multi-Drug Resistant TB (MDR-TB) Patients* confirmed in LA County 2009-2015



*Includes MDR, pre-XDR, and XDR patients




Multi-Drug Resistant TB (MDR-TB) Patients: LA County 2009-2015




LA County TB Patients. Data exclude Pasadena and Long Beach TB patients.
Based on TRIMS data, updated 2/8/16. Data are provisional and subject to change.
*Includes only confirmed (counted) patients

MDR-TB Cases by MDR Infection History




TB Patients by Type of Therapy Administration*: LA County 2011-2015




LA County TB Patients. Data exclude Pasadena and Long Beach TB patients.
Based on TRIMS data, updated 2/8/16. Data are provisional and subject to change.
*Direct Observed Therapy (DOT); Self Administered Therapy (SAT).




TB mortality in LA County vs CA, vs US



Deaths Among TB Patients: LA County 2012-2014




Deaths in Persons with Tuberculosis: LA County 2011-2014



*Tx=Treatment. Cause of death not available at this time.
 LA County TB Patients. Data exclude Pasadena and Long Beach TB patients.
 Based on TRIMS data, updated 2/8/16. Data are provisional and subject to change.



Mortality by Age Group



Case Closure

LA County, 2009 – 2013, N=3,163




POPULATIONS OF SPECIAL INTEREST


HOMELESS




TB Patients Experiencing Homelessness: LA County 2000-2015




Homeless TB Patients by Age: LA County 2015




Homeless TB Patients by Sex: LA County 2015




Homeless TB Patients by Race/Ethnicity: LA County 2015



Homeless TB Patients by Country of Birth: LA County 2015



Risk Factors and Mortality among Homeless TB Patients: LA County 2015



Homeless TB Patients Hospital source: LA County 2007-2015


	Homeless Cases				All
	LAC/USC		Non LAC/USC		
	<u>n</u>	<u>%</u>	<u>n</u>	<u>%</u>	<u>n</u>
All	163	38%	258	61%	421
-					
2007	23	43%	30	56%	53
2008	26	44%	32	55%	58
2009	14	37%	23	62%	37
2010	5	14%	29	85%	34
2011	22	41%	31	58%	53
2012	13	33%	26	66%	39
2013	28	42%	38	57%	66
2014	17	45%	20	54%	37
2015	15	34%	29	65%	44



TB Outbreak #1 Update: Epi-Curve

Genotypic cluster G11610:

“Large Homeless Outbreak”





LA County TB Patients. Data exclude Pasadena and Long Beach TB patients.
 Based on TRIMS and data from the Genotype Cluster and Investigation Assessment Unit, TBCU, updated 2/17/16.
 Data are provisional and subject to change. Patients are defined as "homeless" per RVCT definition.



TB Outbreak Update: Demographics of Homeless patients in Genotypic cluster G11610: LA County 2007-2015

**Outbreak of TB in the Homeless,
2007-2015: Age Distribution**




LA County TB Patients. Data exclude Pasadena and Long Beach TB patients.
Based on TRIMS and data from the Genotype Cluster and Investigation Assessment Unit, TBCP, updated 2/17/16.
Data are provisional and subject to change. Patients are defined as "homeless" per RVCT definition.




TB Outbreak Update: Demographics of Homeless patients in Genotypic cluster G11610: LA County 2007-2015

Country of Birth



Race/ethnicity



TB Screening and Treatment among Private Providers Serving Homeless Populations

- Capturing TB screening records for TB clearance
- Encouraging 3HP use
- TBCP weekly screening @ Pathways
- Coming soon: CHS run onsite Pathways to Home clinic



LOS ANGELES CHRISTIAN
HEALTH CENTERS
"SERVING ALL, WITH OPEN HEARTS"

Logo sources:

URM - <http://urm.org/> CCCHC - <http://centralcityhealth.org/>

JWCH - <http://jwchinstitute.org/> LACHC - <http://www.lachc.com/>

SJWCFC - <http://www.wellchild.org/>



TB Screening and Treatment among Private Providers Serving Homeless Populations

- Since 2013, screened 3186
- 408 lost
- 307 TBI+
- 92 started tx
- 57 (62%) completed



LOS ANGELES CHRISTIAN
HEALTH CENTERS
"SERVING ALL, WITH OPEN HEARTS"

Logo sources:

URM - <http://urm.org/> CCCHC - <http://centralcityhealth.org/>

JWCH - <http://jwchinstitute.org/> LACHC - <http://www.lachc.com/>

SJWCFC - <http://www.wellchild.org/>



Homeless 3HP Treatment Status– CHS and LACHC As of February 9, 2016

Status	Homeless				Total	% among Patients with Closure
	Yes	%	No	%		
Initiated Treatment	240		46		286	
Pending Closure	22	9%	6	13%	28	10%
Completed 3HP Treatment	157	65%	37	80%	194	68%
Stopped, Medical Reason*	9	4%	2	4%	11	4%
Stopped, Other Reason**	52	22%	1	2%	53	18%

*Includes 8 patients stopping due to AE: 4 elevated LFT's, 1 nausea/vomiting, 1 dry mouth, 1 rash, 1 other
 The overwhelming majority of these patients were treated by CHS, and were homeless.
 14 patients were identified as Contacts.
 9 patients were identified as B-notifications.
 All were administered by DOT.
 ** Other reason includes AMA, lost, moved, or previous treatment



SPECIAL PROJECTS



Outbreak #1 Homeless Outbreak

Contact Investigation at Homeless Shelter

- Large number of homeless cluster patients have been housed at one large shelter in LA County
- Ongoing transmission has been suspected
- CDC conducted a large outbreak investigation, (known as Epi Aid) in 2013, the methods of which can be applied to prioritization of contacts in ongoing contact investigations (CIs)



2013 EpiAid Investigation at Homeless Shelter

2013 CDC Epi Aid Contact Prioritization		
Tier	Definition	Identified (n)
I	HIV+, or >120 days of exposure	2,474
II	30-120 days of exposure AND last exposed AFTER January 1, 2012	1,690
III	30-120 days of exposure AND last exposed BEFORE January 1, 2012	1,267
IV	1-29 days of exposure	12,226
Total		17,657

Note: Included contacts to 45 cases with exposure in various shelters



2015 Contact Investigation at Homeless Shelter: Patient Information

- 3 homeless TB patients (all smear 4+/cavitary) presented in early 2015
- All were at the same shelter during their infectious periods –their exposure periods were consecutive and non-overlapping between 10/2014 and 2/2015
- Performed one combined CI for contacts to all three patients at the shelter using CDC Epi Aid prioritization methodology



2015 Contact Investigation at Homeless Shelter: Contact Prioritization

Tier	Definition	Identified (n)
I	≥ 60 total days of exposure, OR ≥ 50 days of exposure to case #1*	173
II	Not in Tier I, and total days of exposure ≥ 30 and < 60	249
III	Total days of exposure ≥ 1 and < 30	871
Total**		1,293

*A holdover from initial CI assessment of the three patients separately


**Excluding n=2 contacts Tier IV (0 days exposure)



2015 Contact Investigation at Homeless Shelter: Contact Prioritization

Overall Screening Results		
Tier	Evaluated (n)	QFT+ (n (%))
I	68	26 (38%)
II	18	4 (22%)
III	24	6 (25%)
Total	110	36 (33%)

Note: 5 Tier I and 1 Tier III contacts lost to X-ray follow up – LTBI confirmed rates are 31% and 19% for Tier I and Tier III, respectively.




*A holdover from initial CI assessment of the three patients separately

**Excluding n=2 contacts Tier IV (0 days exposure)




Contact Investigation at Homeless Shelter: Screening Results

Total Screened Contacts by Exposure Days



Contact Investigation at Homeless Shelter: Screened Tier I Contacts (N=68) according to prior 2013 CDC EPI Aid Tier Status



Contact Investigation at Homeless Shelter: Summary

- LTBI data corroborate findings of 2013 Epi Aid and support this prioritization method
- Use of shelter registration data to calculate smear positive/cavitary exposure nights is useful for prioritization at this shelter
- Data show importance of contact prioritization and investigation based on recognition of the ongoing outbreak, not by individual patients




Outbreak #2

Grupo Cluster: Genotype G10161

- What are Grupos?
 - Community-based alcohol rehabilitation centers that follow a faith-based alcoholic recovery program
 - Population consists of clients that are predominantly from Central or South America
 - Provide cooperative living arrangements and meeting space for recovering drug and alcohol users
 - Sites offer dormitory and communal housing and are supported by their large network of members



Grupo Cluster: Locations in LA County



Grupo Cluster: Demographics

- 26 TB patients in LAC (2007-2015)
 - 6 confirmed TB patients in 2015
 - 23 are non-U.S. born:
 - Mexico (10)
 - El Salvador (8)
 - Guatemala (3)
 - Honduras (1)
 - Argentina (1)
 - Main risk factor: exposure to one or more Grupos
 - 58% of patients have some degree of homelessness



Grupo Cluster: Demographics

Comparison between LA County and U.S.

Characteristic	Los Angeles N=26	Rest of U.S. N=8
Race/Ethnicity		
Hispanic	26 (100.0%)	7 (87.5%)
Asian	0 (0.0)	1 (12.5%)
Gender		
Male	21 (91.3%)	6 (75.0%)
Female	2 (8.7%)	2 (25.0%)
Birth		
US Born	2 (7.7%)	0 (0.0%)
non-U.S. Born	23 (88.5%)	8 (100.0%)
Unknown	1 (3.8%)	




Grupo Cluster: Risk Factors: LA County

	Grupo Cluster-LAC	
	<u>n</u>	<u>%</u>
Selected Risk Factors	26	
Homelessness		
Yes	15	58.0
HIV		
Yes	3	12.0
Diabetes		
Yes	6	23.1
Injecting Drug Use		
Yes	2	8.0
Non-injecting Drug Use		
Yes	7	27.0
Alcohol Use		
Yes	22	85.0



Grupo Cluster: Epi Curve



Grupo Cluster: Cases in G10161/G10167, PCR00556 (n=20) with Grupo History During Exposure Period: LA County 2007-2015

Tier I: Red solid triangle

1. Grupos with cases that are cavitory and sputum smear +
2. A Grupo that has two or more cases with one of these cases being cavitory and sputum smear +

Tier II: Blue solid triangle

1. Grupos with cases that are non-cavitory and sputum smear +, or
2. Any case that is cavitory and sputum smear negative

Tier III: Green solid triangle

1. Grupos with cases that are non-cavitory and sputum smear –





Image 1: 20 of the 26 cases in G10161/G10167, PCR00556 had Grupo history during exposure period identified during patient interviews

Source: Genotype Cluster and Investigation Assessment Unit, TBPCP, NodeXL.

Coming soon:

- NAAT performance results
- Analysis of TBI Treatment in Contacts
- Implementation and Evaluation of Automated VDOT
- Mortality project continues.....



Final Thoughts

How can I use
epi data to
communicate
risk?

Am I under-
reporting or
misclassifying
data?

How can I use
data to prioritize
and evaluate my
work?



**THANK YOU!
ANY QUESTIONS?**

